

PROCESSED AT :  
Thyrocare



Corporate office : Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400 703  
☎ 022 - 3090 0000 / 6712 3400 📞 9870666333 ✉ wellness@thyrocare.com 🌐 www.thyrocare.com

REPORT

NAME : XXXXXXXXXXXXXXXXXXXX  
REF. BY : XXXXXXXXXXXXXXXXXXXX  
TEST ASKED : CRP

SAMPLE COLLECTED AT :  
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
C-REACTIVE PROTEIN (CRP)	IMMUNOTURBIDIMETRY	23.78	mg/L

Reference Range : (mg/L)

Acute phase determination : < 5 mg/L

Clinical Significance:

It's a protein present in the sera of acutely ill patients that bound cell wall C-polysaccharide of streptococcus pneumoniae and agglutinates the organisms.

CRP is one of the strongest acute -phase reactants, with plasma concentrations rising up after myocardial infarction, stress, trauma, infection, inflammation, surgery, or neoplastic proliferation.

Concentrations >5 to 10mg/L suggest the presence of an infection or inflammatory process. Concentrations are generally higher in bacterial than viral infection. The increase in peak is proportional to tissue damage. Determination of CRP is clinically useful to screen activity of inflammatory diseases such as rheumatoid arthritis; SLE; Leukemia; after surgery; to detect rejection in renal allograft recipients; to detect neonatal septicemia and meningitis. However, its is a nonspecific marker and cannot be interpreted without other

Please correlate with clinical conditions.

~~ End of report ~~

Sample Collected on (SCT) : Sample collection time  
Sample Received on (SRT) : Sample receiving time at Lab  
Report Released on (RRT) : Report release time  
Sample Type : SERUM  
Labcode :  
Barcode :

Doctor 1 Sign

Doctor 2 Sign