



Tests you can trust

Name : XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

Date : XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

Test Asked : Senior Citizen Profile Male

Report Status: Complete Report



First National Diagnostic Chain to have
100% of its Labs with NABL Accreditation[#]



98% Reports
released within
06 Hours
of sample reaching the lab*



9 out of 10 Doctors Trust
that Thyrocare
reports are
Accurate & Reliable*



1200+
Tests & Profiles



Temperature-
Controlled
Sample Logistics



Unique Barcode
Tracking



Fully Automated
Machines Inspected
Daily



Abnormal Values
Re-Checked Twice



Reports Verified By Expert
MD Pathologists Stationed
at Every Lab



Your reports are digitally verifiable

Scan the QR code inside the report to check authenticity
of reported values

QR code will remain active for 30 days from report release date

Accredited by



NABL From 2005



ISO 9001: 2015 - From 2015



CAP From 2007

PROCESSED AT :

 Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400 703  9870666333  wellness@thyrocare.com

First National Diagnostic Chain to have 100% of its Labs with NABL Accreditation[#]

NAME : XXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXX
TEST ASKED : SENIOR CITIZEN PROFILE MALE

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXX

Report Availability Summary

Note: Please refer to the table below for status of your tests.

 **22** Ready  **0** Ready with Cancellation  **0** Processing  **0** Cancelled in Lab

TEST DETAILS

REPORT STATUS

SENIOR CITIZEN PROFILE MALE

Ready 

FASTING BLOOD SUGAR (GLUCOSE)

Ready 

CARCINO EMBRYONIC ANTIGEN (CEA)

Ready 

CHLORIDE

Ready 

25-OH VITAMIN D (TOTAL)

Ready 

FERRITIN

Ready 

HIGH SENSITIVITY C-REACTIVE PROTEIN (HS-CRP)

Ready 

LIPASE

Ready 

Lipoprotein (a) [Lp(a)]

Ready 

PROSTATE SPECIFIC ANTIGEN (PSA)

Ready 

SODIUM

Ready 

VITAMIN B-12

Ready 

COMPLETE URINE ANALYSIS

Ready 

HBA PROFILE

Ready 

HEMOGRAM - 6 PART (DIFF)

Ready 

LIVER FUNCTION TESTS

Ready 

ELEMENTS 22 (TOXIC AND NUTRIENTS)

Ready 

IRON DEFICIENCY PROFILE

Ready 

KIDPRO

Ready 

LIPID PROFILE

Ready 

T3-T4-USTSH

Ready 

PROCESSED AT :



Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400 703 | 9870666333 | wellness@thyrocare.com

First National Diagnostic Chain to have 100% of its Labs with NABL Accreditation#

NAME : XXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXX
TEST ASKED : SENIOR CITIZEN PROFILE MALE
SAMPLE COLLECTED AT : XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

Report Availability Summary

Note: Please refer to the table below for status of your tests.

22 Ready 0 Ready with Cancellation 0 Processing 0 Cancelled in Lab

TEST DETAILS	REPORT STATUS
APOLIPROTEIN RATIO	Ready
AMYLASE	Ready

PROCESSED AT :



Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400 703 | 9870666333 | wellness@thyrocare.com

First National Diagnostic Chain to have 100% of its Labs with NABL Accreditation[#]

NAME : XXXXXXXXXXXXXXXXXXXX SAMPLE COLLECTED AT :
 REF. BY : XXXXXXXXXXXXXXXXXXXX XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX
 TEST ASKED : SENIOR CITIZEN PROFILE MALE

Summary Report

Tests outside reference range

TEST NAME	OBSERVED VALUE	UNITS	Bio. Ref. Interval.
COMPLETE HEMOGRAM			
HEMATOCRIT(PCV)	33.6	%	40.0-50.0
HEMOGLOBIN	10.4	g/dL	13.0-17.0
LYMPHOCYTE	17.4	%	20-40
MEAN CORP. HEMO. CONC(MCHC)	31	g/dL	31.5-34.5
MEAN CORPUSCULAR HEMOGLOBIN(MCH)	26.8	pq	27.0-32.0
RED CELL DISTRIBUTION WIDTH (RDW-CV)	15.4	%	11.6-14
RED CELL DISTRIBUTION WIDTH - SD(RDW-SD)	48.9	fL	39-46
TOTAL RBC	3.88	X 10 ⁶ /μL	4.5-5.5
LIPID			
LDL / HDL RATIO	0.9	Ratio	1.5-3.5
TC/ HDL CHOLESTEROL RATIO	2.2	Ratio	3 - 5
RENAL			
CREATININE - SERUM	0.7	mg/dL	0.72-1.18

Disclaimer: The above listed is the summary of the parameters with values outside the BRI. For detailed report values, parameter correlation and clinical interpretation, kindly refer to the same in subsequent pages.

PROCESSED AT :



Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400 703 | 9870666333 | wellness@thyrocare.com

First National Diagnostic Chain to have 100% of its Labs with NABL Accreditation[#]

NAME : XXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXX
TEST ASKED : SENIOR CITIZEN PROFILE MALE

SAMPLE COLLECTED AT :

XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
CARCINO EMBRYONIC ANTIGEN (CEA)	C.L.I.A	< 1.25	ng/mL
Bio. Ref. Interval. :-			

Non-Smokers : < 2.50 ng/mL

Smokers : < 5.00 ng/mL

Clinical Significance :

CEA is often used to monitor patients with cancers of the gastrointestinal tract (GI). Increased CEA levels can indicate some Non-Cancer related conditions, Such as some forms of inflammation, Cirrhosis, and Peptic Ulcer. Also, Smokers tend to have Higher CEA levels than Non-Smokers. When cancer spreads to other organs, CEA levels rise and may be present in other types of bodily fluids besides blood.

For Diagnostic Purpose, Results should always be assessed in Conjunction with the patients medical history, clinical examination and other findings.

Specifications:

Precision: Intra Assay (%CV): 3.6 %, Inter Assay (%CV): 4.1 %; Sensitivity: 0.5 ng/ml

Kit Validation References:

Statland Be, Winkel P. Neoplasia. In: Kaplan LA, Resc AJ, Editors. Clinical Chemistry, Theory, Analysis and Correlation. 2nd Ed. St. Louis: Cv Mosby, 1989.p 734-5.

Please correlate with clinical conditions.

Method:- FULLY AUTOMATED TWO STEP SANDWICH IMMUNOASSAY

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign

Doctor 2 Sign

PROCESSED AT :



Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400 703 | 9870666333 | wellness@thyrocare.com

First National Diagnostic Chain to have 100% of its Labs with NABL Accreditation[#]

NAME : XXXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXXX
TEST ASKED : SENIOR CITIZEN PROFILE MALE

SAMPLE COLLECTED AT :
 XX

TEST NAME	TECHNOLOGY	VALUE	UNITS
25-OH VITAMIN D (TOTAL)	C.L.I.A	< 65	ng/mL
Bio. Ref. Interval. :-			

DEFICIENCY : <20 ng/ml || INSUFFICIENCY : 20-<30 ng/ml
 SUFFICIENCY : 30-100 ng/ml || TOXICITY : >100 ng/ml

Clinical Significance:

Vitamin D is a fat soluble vitamin that has been known to help the body absorb and retain calcium and phosphorous; both are critical for building bone health. Decrease in vitamin D total levels indicate inadequate exposure of sunlight, dietary deficiency, nephrotic syndrome. Increase in vitamin D total levels indicate Vitamin D intoxication.

Specifications: Precision: Intra assay (%CV):5.3%, Inter assay (%CV):11.9% ; Sensitivity:3.2 ng/ml.

Kit Validation Reference: Holick MF. Vitamin D Deficiency. N Engl J Med. 2007;357:266-81.

Please correlate with clinical conditions.

Method:- Fully Automated Chemi Luminescent Immuno Assay

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign

Doctor 2 Sign

PROCESSED AT :



Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400 703 | 9870666333 | wellness@thyrocare.com

First National Diagnostic Chain to have 100% of its Labs with NABL Accreditation[#]

NAME : XXXXXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXXXXX
TEST ASKED : SENIOR CITIZEN PROFILE MALE

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
APOLIPOPROTEIN - A1 (APO-A1)	IMMUNOTURBIDIMETR Y	< 119	mg/dL
Bio. Ref. Interval. : Male : 86 - 152 Female : 94 - 162 Method : FULLY AUTOMATED RATE IMMUNOTURBIDIMETRY - BECKMAN COULTER			
APOLIPOPROTEIN - B (APO-B)	IMMUNOTURBIDIMETR Y	< 101	mg/dL
Bio. Ref. Interval. : Male : 56 - 145 Female : 53 - 138 Method : FULLY AUTOMATED RATE IMMUNOTURBIDIMETRY - BECKMAN COULTER			
APO B / APO A1 RATIO (APO B/A1)	CALCULATED	0.8	Ratio
Bio. Ref. Interval. : Male : 0.40 - 1.26 Female : 0.38 - 1.14 Clinical Significance : • Apolipoprotein B is a more potent and independent predictor of Coronary artery disease (CAD) than LDL Cholesterol. • Apolipoprotein A1 is one of the apoproteins of HDL and is inversely related to risk of CAD. • The Apolipoprotein studies help in monitoring risk of restenosis in patients with myocardial infarction, Coronary bypass surgery etc. • An increased ratio of Apo B to A1 beyond the defined normal range is indicative of CAD risk. • All results have to be interpreted in Conjunction with clinical history and other findings. Method : Derived from serum Apo A1 and Apo B values			

Please correlate with clinical conditions.

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign

Doctor 2 Sign

PROCESSED AT :



Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400 703 | 9870666333 | wellness@thyrocare.com

First National Diagnostic Chain to have 100% of its Labs with NABL Accreditation[#]

NAME : XXXXXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXXXXX
TEST ASKED : SENIOR CITIZEN PROFILE MALE

SAMPLE COLLECTED AT :

XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
HIGH SENSITIVITY C-REACTIVE PROTEIN (HS-CRP) Bio. Ref. Interval. :-	IMMUNOTURBIDIMETRY	< 1.5	mg/L

< 1.00 - Low Risk
1.00 - 3.00 - Average Risk
>3.00 - 10.00 - High Risk
> 10.00 - Possibly due to Non-Cardiac Inflammation

Disclaimer: Persistent unexplained elevation of HSCRP >10 should be evaluated for non-cardiovascular etiologies such as infection, active arthritis or concurrent illness.

Clinical significance:

High sensitivity C- reactive Protein (HSCRP) can be used as an independent risk marker for the identification of Individuals at risk for future cardiovascular Disease. A coronary artery disease risk assessment should be based on the average of two hs-CRP tests, ideally taken two weeks apart.

Kit Validation Reference:

- 1.Clinical management of laboratory data in medical practice 2003-2004, 207(2003).
- 2.Tietz : Textbook of Clinical Chemistry and Molecular diagnostics :Second edition :Chapter 47:Page no.1507- 1508.

Please correlate with clinical conditions.

Method:- FULLY AUTOMATED LATEX AGGLUTINATION - BECKMAN COULTER

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign

Doctor 2 Sign

PROCESSED AT :



Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400 703 | 9870666333 | wellness@thyrocare.com

First National Diagnostic Chain to have 100% of its Labs with NABL Accreditation[#]

NAME : XXXXXXXXXXXXXXXXXXXX
 REF. BY : XXXXXXXXXXXXXXXXXXXX
 TEST ASKED : SENIOR CITIZEN PROFILE MALE
 SAMPLE COLLECTED AT : XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
FERRITIN Bio. Ref. Interval. : Men: 22-322 ng/ml Women: 10-291 ng/ml Method : Fully Automated Bidirectionally Interfaced Chemi Luminescent Immuno Assay	C.L.I.A	< 215	ng/mL
IRON Bio. Ref. Interval. : Male : 65 - 175 Female : 50 - 170 Method : Ferrozine method without deproteinization	PHOTOMETRY	< 120	µg/dL
TOTAL IRON BINDING CAPACITY (TIBC) Bio. Ref. Interval. : Male: 225 - 535 µg/dl Female: 215 - 535 µg/dl Method : Spectrophotometric Assay	PHOTOMETRY	< 380	µg/dL
% TRANSFERRIN SATURATION Bio. Ref. Interval. : 13 - 45 Method : Derived from IRON and TIBC values	CALCULATED	32	%
UNSAT. IRON-BINDING CAPACITY (UIBC) Bio. Ref. Interval. : 162 - 368 Method : SPECTROPHOTOMETRIC ASSAY	PHOTOMETRY	< 265	µg/dL

Please correlate with clinical conditions.

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign Doctor 2 Sign

PROCESSED AT :



Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400 703 | 9870666333 | wellness@thyrocare.com

First National Diagnostic Chain to have 100% of its Labs with NABL Accreditation[#]

NAME : XXXXXXXXXXXXXXXX
 REF. BY : XXXXXXXXXXXXXXXX
 TEST ASKED : SENIOR CITIZEN PROFILE MALE
 SAMPLE COLLECTED AT : XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
VITAMIN B-12	C.L.I.A	< 484	pg/mL
Bio. Ref. Interval. :-			

Normal : 211 - 911 pg/ml

Clinical significance :

Vitamin B12 or cyanocobalamin, is a complex corrinoid compound found exclusively from animal dietary sources, such as meat, eggs and milk. It is critical in normal DNA synthesis, which in turn affects erythrocyte maturation and in the formation of myelin sheath. Vitamin-B12 is used to find out neurological abnormalities and impaired DNA synthesis associated with macrocytic anemias. For diagnostic purpose, results should always be assessed in conjunction with the patients medical history, clinical examination and other findings.

Specifications: Intra assay (%CV):5.0%, Inter assay (%CV):9.2 %;Sensitivity:45 pg/ml

Kit Validation reference:

Chen IW, Sperling MI, Heminger LA. Vitamin B12. In: Pesce AJ, Kaplan LA, eds. Methods in Clinical Chemistry. St. Louis: CV Mosby; 1987:569-73.

Please correlate with clinical conditions.

Method:- COMPETITIVE CHEMI LUMINESCENT IMMUNO ASSAY

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign

Doctor 2 Sign

PROCESSED AT :



Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400 703 | 9870666333 | wellness@thyrocare.com

First National Diagnostic Chain to have 100% of its Labs with NABL Accreditation[#]

NAME : XXXXXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXXXXX
TEST ASKED : SENIOR CITIZEN PROFILE MALE
SAMPLE COLLECTED AT : XX

TEST NAME	TECHNOLOGY	VALUE	UNITS
Lipoprotein (a) [Lp(a)] Bio. Ref. Interval. :-	IMMUNOTURBIDIMETRY	< 14.95	mg/dL

Adults : < 30.0 mg/dl

Clinical Significance:

Determination of LPA may be useful to guide management of individuals with a family history of CHD or with existing disease. The levels of LPA in the blood depends on genetic factors; The range of variation in a population is relatively large and hence for diagnostic purpose, results should always be assessed in conjunction with the patient's medical history, clinical examination and other findings.

Specifications:

Precision %CV :- Intra assay %CV- 4.55% , Inter assay %CV-0.86 %

Kit Validation Reference:

Tietz NW, Clinical Guide to Laboratory Tests Philadelphia WB. Saunders 1995 : 442-444

Please correlate with clinical conditions.

Method:- LATEX ENHANCED IMMUNOTURBIDIMETRY

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign

Doctor 2 Sign

Page : 7 of 22

PROCESSED AT :



Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400 703 | 9870666333 | wellness@thyrocare.com

First National Diagnostic Chain to have 100% of its Labs with NABL Accreditation[#]

NAME : XXXXXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXXXXX
TEST ASKED : SENIOR CITIZEN PROFILE MALE
SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
PROSTATE SPECIFIC ANTIGEN (PSA) Bio. Ref. Interval. :-	C.L.I.A	< 0.7	ng/mL

Normal : < 4.00 ng/ml
Border line : 4.01 to 10.00 ng/ml

Clinical Significance:

Elevated levels of PSA are associated with prostate cancer, but may also be seen with prostatitis (Inflammation of the prostate) and benign prostatic hyperplasia (BPH). PSA test done along with free PSA provides additional information. Studies have suggested that the percentage of free PSA in total PSA is lower in patients with prostate cancer than those with benign prostate hyperplasia.

Specification:

Precision: Intra assay (%CV): 4.38%, Inter assay (%CV): 4.67%; Sensitivity: 0.01 ng/ml

Kit validation references:

Wang MC, Valenzuela LA, Murphy GP, and Chu TM. Purification of a human prostate-specific antigen. Invest. Urol. 1979; 17: 159

Please correlate with clinical conditions.

Method:- TWO SITE SANDWICH IMMUNOASSAY

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign

Doctor 2 Sign

Page : 8 of 22

First National Diagnostic Chain to have 100% of its Labs with NABL Accreditation[#]

NAME : XXXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXXX
TEST ASKED : SENIOR CITIZEN PROFILE MALE

SAMPLE COLLECTED AT :
 XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
AMYLASE	PHOTOMETRY	< 64	U/L

Bio. Ref. Interval. :-

Adults : 28-100 U/L

Interpretation:

Lipemic Sera (Hypertriglyceridemia) may contain inhibitors, Which falsely depress results. About 20% of patients with Acute Pancreatitis have abnormal lipids. Normal serum amylase may occur in Pancreatitis, Especially relapsing and chronic pancreatitis. Moderate increases may be reported in normal pregnancy.

Clinical Significance:

Causes of high Serum Amylase include Acute Pancreatitis, Pancreatic Pseudocyst, Pancreatic Ascites, Pancreatic Abscess, Neoplasm in or adjacent to Pancreas, Trauma to Pancreas, and common Duct Stones. Nonpancreatic Causes include inflammatory salivary lesions (Eg, Mumps), Perforated Peptic Ulcer, Intestinal Obstruction, Biliary Tract Disease, Peritonitis, Acute Appendicitis, Diabetic Ketoacidosis, and Extrapneumatic Carcinomas. Amylase levels more than 25-fold the upper limit of normal are often found when metastatic tumors produce Ectopic Amylase.

Specifications:

Precision: Intra assay (%CV): 2.82, Inter assay (%CV): 2.49, Sensitivity: 10.9 U/L.

Kit Validation References:

Rauscher, E., et coll., Fresenius Z. Analyt. Chem. 324 (1986) 304-305.

Please correlate with clinical conditions.

Method:- ENZYMATIC COLORIMETRIC TEST

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign

Doctor 2 Sign

PROCESSED AT :



Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400 703 | 9870666333 | wellness@thyrocare.com

First National Diagnostic Chain to have 100% of its Labs with NABL Accreditation[#]

NAME : XXXXXXXXXXXXXXXXXXXX
 REF. BY : XXXXXXXXXXXXXXXXXXXX
 TEST ASKED : SENIOR CITIZEN PROFILE MALE
 SAMPLE COLLECTED AT : XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
LIPASE	PHOTOMETRY	< 28.45	U/L

Bio. Ref. Interval. :-

Adults : 5.6 - 51.3 U/L

Interpretation:

For diagnostic purposes, the results should always be assessed in conjunction with the patient's medical history, clinical examination and other findings like serum amylase. Serum Lipase is usually normal in patients with elevated serum amylase, having peptic ulcer, salivary adenitis, inflammatory bowel disease, intestinal obstruction, and macroamylasemia. Lipemic sera may interfere with results.

Clinical Significance:

High serum Lipase is a specific marker for pancreatitis; after acute pancreatitis the Lipase activity increases within 4-8 hours, reaches a peak after 24 hours and decreases after 8 to 14 days. However, there is no correlation between the Lipase activity determined in serum and the extent of damage to the pancreas.

Specifications:

Precision: Intra assay (%CV): 3.35, Inter assay (%CV): 2.46, Sensitivity: 3.5 U/L.

Kit Validation References:

Tietz Nw Et Al. Lipase In Serum - The Elusive Enzyme: An Overview. Clin Chem 1993; 39:746-756.

Please correlate with clinical conditions.

Method:- ENZYMATIC COLORIMETRIC ASSAY

Sample Collected on (SCT) : Sample collection time
 Sample Received on (SRT) : Sample receiving time at Lab
 Report Released on (RRT) : Report release time
 Sample Type : SERUM
 Labcode :
 Barcode :

Doctor 1 Sign

Doctor 2 Sign

First National Diagnostic Chain to have 100% of its Labs with NABL Accreditation[#]

NAME : XXXXXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXXXXX
TEST ASKED : SENIOR CITIZEN PROFILE MALE

SAMPLE COLLECTED AT :

XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS	Bio. Ref. Interval.
TOTAL CHOLESTEROL	PHOTOMETRY	123	mg/dL	< 200
HDL CHOLESTEROL - DIRECT	PHOTOMETRY	56	mg/dL	40-60
LDL CHOLESTEROL - DIRECT	PHOTOMETRY	53.5	mg/dL	< 100
TRIGLYCERIDES	PHOTOMETRY	120	mg/dL	< 150
TC/ HDL CHOLESTEROL RATIO	CALCULATED	2.2	Ratio	3 - 5
TRIG / HDL RATIO	CALCULATED	2.12	Ratio	< 3.12
LDL / HDL RATIO	CALCULATED	0.9	Ratio	1.5-3.5
HDL / LDL RATIO	CALCULATED	1.05	Ratio	> 0.40
NON-HDL CHOLESTEROL	CALCULATED	66.2	mg/dL	< 160
VLDL CHOLESTEROL	CALCULATED	23.94	mg/dL	5 - 40

Please correlate with clinical conditions.

Method :

CHOL - Cholesterol Oxidase, Esterase, Peroxidase
HCHO - Direct Enzymatic Colorimetric
LDL - Direct Measure
TRIG - Enzymatic, End Point
TC/H - Derived from serum Cholesterol and Hdl values
TRI/H - Derived from TRIG and HDL Values
LDL/ - Derived from serum HDL and LDL Values
HD/LD - Derived from HDL and LDL values.
NHDH - Derived from serum Cholesterol and HDL values
VLDL - Derived from serum Triglyceride values

***REFERENCE RANGES AS PER NCEP ATP III GUIDELINES:**

TOTAL CHOLESTEROL	(mg/dl)	HDL	(mg/dl)	LDL	(mg/dl)	TRIGLYCERIDES	(mg/dl)
DESIRABLE	<200	LOW	<40	OPTIMAL	<100	NORMAL	<150
BORDERLINE HIGH	200-239	HIGH	>60	NEAR OPTIMAL	100-129	BORDERLINE HIGH	150-199
HIGH	>240			BORDERLINE HIGH	130-159	HIGH	200-499
				HIGH	160-189	VERY HIGH	>500
				VERY HIGH	>190		

Alert !!! 10-12 hours fasting is mandatory for lipid parameters. If not, values might fluctuate.

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labode :
Barcode :

Doctor 1 Sign

Doctor 2 Sign

First National Diagnostic Chain to have 100% of its Labs with NABL Accreditation[#]

NAME : XXXXXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXXXXX
TEST ASKED : SENIOR CITIZEN PROFILE MALE

SAMPLE COLLECTED AT :

XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS	Bio. Ref. Interval.
ALKALINE PHOSPHATASE	PHOTOMETRY	< 87	U/L	45-129
BILIRUBIN - TOTAL	PHOTOMETRY	< 0.75	mg/dL	0.3-1.2
BILIRUBIN -DIRECT	PHOTOMETRY	< 0.15	mg/dL	< 0.3
BILIRUBIN (INDIRECT)	CALCULATED	0.6	mg/dL	0-0.9
GAMMA GLUTAMYL TRANSFERASE (GGT)	PHOTOMETRY	< 27.45	U/L	< 55
ASPARTATE AMINOTRANSFERASE (SGOT)	PHOTOMETRY	< 17.5	U/L	< 35
ALANINE TRANSAMINASE (SGPT)	PHOTOMETRY	< 22.5	U/L	< 45
SGOT / SGPT RATIO	CALCULATED	0.78	Ratio	< 2
PROTEIN - TOTAL	PHOTOMETRY	< 6.95	gm/dL	5.7-8.2
ALBUMIN - SERUM	PHOTOMETRY	< 4	gm/dL	3.2-4.8
SERUM GLOBULIN	CALCULATED	< 2.95	gm/dL	2.5-3.4
SERUM ALB/GLOBULIN RATIO	CALCULATED	1.36	Ratio	0.9 - 2

Please correlate with clinical conditions.

Method :

ALKP - Modified IFCC method
 BILT - Vanadate Oxidation
 BILD - Vanadate Oxidation
 BILI - Derived from serum Total and Direct Bilirubin values
 GGT - Modified IFCC method
 SGOT - IFCC* Without Pyridoxal Phosphate Activation
 SGPT - IFCC* Without Pyridoxal Phosphate Activation
 OT/PT - Derived from SGOT and SGPT values.
 PROT - Biuret Method
 SALB - Albumin Bcg¹ method (Colorimetric Assay Endpoint)
 SEGB - DERIVED FROM SERUM ALBUMIN AND PROTEIN VALUES
 A/GR - Derived from serum Albumin and Protein values

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign

Doctor 2 Sign

PROCESSED AT :



Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400 703 | 9870666333 | wellness@thyrocare.com

First National Diagnostic Chain to have 100% of its Labs with NABL Accreditation[#]

NAME : XXXXXXXXXXXXXXXXXXXX SAMPLE COLLECTED AT :
 REF. BY : XXXXXXXXXXXXXXXXXXXX XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX
 TEST ASKED : SENIOR CITIZEN PROFILE MALE

TEST NAME	TECHNOLOGY	VALUE	UNITS
CHLORIDE Bio. Ref. Interval. : ADULTS: 98-107 MMOL/L	I.S.E - INDIRECT	< 102.5	mmol/L
Clinical Significance : An increased level of blood chloride (called hyperchloremia) usually indicates dehydration, but can also occur with other problems that cause high blood sodium, such as Cushing syndrome or kidney disease. Hyperchloremia also occurs when too much base is lost from the body (producing metabolic acidosis) or when a person hyperventilates (causing respiratory alkalosis). A decreased level of blood chloride (called hypochloremia) occurs with any disorder that causes low blood sodium. Hypochloremia also occurs with congestive heart failure, prolonged vomiting or gastric suction, Addison disease, emphysema or other chronic lung diseases (causing respiratory acidosis), and with loss of acid from the body (called metabolic alkalosis). Method : ION SELECTIVE ELECTRODE - INDIRECT			
SODIUM Bio. Ref. Interval. : Adults: 136-145 mmol/l Method : ION SELECTIVE ELECTRODE - INDIRECT	I.S.E - INDIRECT	< 140.5	mmol/L
Please correlate with clinical conditions.			

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign

Doctor 2 Sign

NAME : XXXXXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXXXXX
TEST ASKED : SENIOR CITIZEN PROFILE MALE

SAMPLE COLLECTED AT :

XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS	Bio. Ref. Interval.
BLOOD UREA NITROGEN (BUN)	PHOTOMETRY	< 14.01	mg/dL	7.94 - 20.07
CREATININE - SERUM	PHOTOMETRY	0.7	mg/dL	0.72-1.18
BUN / Sr.CREATININE RATIO	CALCULATED	20.01	Ratio	9:1-23:1
UREA (CALCULATED)	CALCULATED	29.98	mg/dL	Adult : 17-43
UREA / SR.CREATININE RATIO	CALCULATED	42.83	Ratio	< 52
CALCIUM	PHOTOMETRY	< 9.7	mg/dL	8.8-10.6
URIC ACID	PHOTOMETRY	< 5.8	mg/dL	4.2 - 7.3

Please correlate with clinical conditions.

Method :

BUN - Kinetic UV Assay.
SCRE - Creatinine Enzymatic Method
B/CR - Derived from serum Bun and Creatinine values
UREAC - Derived from BUN Value.
UR/CR - Derived from UREA and Sr.Creatinine values.
CALC - Arsenazo III Method, End Point.
URIC - Urilase / Peroxidase Method

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign

Doctor 2 Sign

PROCESSED AT :



Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400 703 | 9870666333 | wellness@thyrocare.com

First National Diagnostic Chain to have 100% of its Labs with NABL Accreditation[#]

NAME : XXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXX
TEST ASKED : SENIOR CITIZEN PROFILE MALE
SAMPLE COLLECTED AT : XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS	Bio. Ref. Interval.
TOTAL TRIIODOTHYRONINE (T3)	C.L.I.A	109	ng/dL	60-200
TOTAL THYROXINE (T4)	C.L.I.A	<8.29	µg/dL	4.5-12
TSH - ULTRASENSITIVE	C.M.I.A	<2.92	µIU/mL	0.35-4.94

The Biological Reference Ranges is specific to the age group. Kindly correlate clinically.

Method :

T3 - Competitive Chemi Luminescent Immuno Assay
T4 - Competitive Chemi Luminescent Immuno Assay
USTSH - Fully Automated Chemi Luminescent Microparticle Immunoassay

Disclaimer : Results should always be interpreted using the reference range provided by the laboratory that performed the test. Different laboratories do tests using different technologies, methods and using different reagents which may cause difference. In reference ranges and hence it is recommended to interpret result with assay specific reference ranges provided in the reports. To diagnose and monitor therapy doses, it is recommended to get tested every time at the same Laboratory.

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign

Doctor 2 Sign

PROCESSED AT :

Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400 703 | 9870666333 | wellness@thyrocare.com

First National Diagnostic Chain to have 100% of its Labs with NABL Accreditation[#]

NAME : XXXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXXX
TEST ASKED : SENIOR CITIZEN PROFILE MALE

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
EST. GLOMERULAR FILTRATION RATE (eGFR)	CALCULATED	133	mL/min/1.73 m2
Bio. Ref. Interval. :-			

> = 90 : Normal
60 - 89 : Mild Decrease
45 - 59 : Mild to Moderate Decrease
30 - 44 : Moderate to Severe Decrease
15 - 29 : Severe Decrease

Clinical Significance

The normal serum creatinine reference interval does not necessarily reflect a normal GFR for a patient. Because mild and moderate kidney injury is poorly inferred from serum creatinine alone. Thus, it is recommended for clinical laboratories to routinely estimate glomerular filtration rate (eGFR), a "gold standard" measurement for assessment of renal function, and report the value when serum creatinine is measured for patients 18 and older, when appropriate and feasible. It cannot be measured easily in clinical practice, instead, GFR is estimated from equations using serum creatinine, age, race and sex. This provides easy to interpret information for the doctor and patient on the degree of renal impairment since it approximately equates to the percentage of kidney function remaining. Application of CKD-EPI equation together with the other diagnostic tools in renal medicine will further improve the detection and management of patients with CKD.

Reference

Levey AS, Stevens LA, Schmid CH, Zhang YL, Castro AF, 3rd, Feldman HI, et al. A new equation to estimate glomerular filtration rate. Ann Intern Med. 2009;150(9):604-12.

Please correlate with clinical conditions.

Method:- CKD-EPI Creatinine Equation

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign

Doctor 2 Sign

PROCESSED AT :



Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400 703 | 9870666333 | wellness@thyrocare.com

First National Diagnostic Chain to have 100% of its Labs with NABL Accreditation[#]

NAME : XXXXXXXXXXXXXXXXXXXX
 REF. BY : XXXXXXXXXXXXXXXXXXXX
 TEST ASKED : COMPLETE URINE ANALYSIS

SAMPLE COLLECTED AT :
 XX

TEST NAME	METHODOLOGY	VALUE	UNITS	Bio. Ref. Interval.
Complete Urinogram				
Physical Examination				
VOLUME	Visual Determination	1	mL	-
COLOUR	Visual Determination	PALE YELLOW	-	Pale Yellow
APPEARANCE	Visual Determination	CLEAR	-	Clear
SPECIFIC GRAVITY	pKa change	< 1.003	-	1.003-1.030
PH	pH indicator	5	-	5-8
Chemical Examination				
URINARY PROTEIN	PEI	ABSENT	mg/dL	Absent
URINARY GLUCOSE	GOD-POD	ABSENT	mg/dL	Absent
URINE KETONE	Nitroprusside	ABSENT	mg/dL	Absent
URINARY BILIRUBIN	Diazo coupling	ABSENT	mg/dL	Absent
UROBILINOGEN	Diazo coupling	Normal	mg/dL	<=0.2
BILE SALT	Hays sulphur	ABSENT	-	Absent
BILE PIGMENT	Ehrlich reaction	ABSENT	-	Absent
URINE BLOOD	Peroxidase reaction	ABSENT	-	Absent
NITRITE	Diazo coupling	ABSENT	-	Absent
LEUCOCYTE ESTERASE	Esterase reaction	ABSENT	-	Absent
Microscopic Examination				
MUCUS	Microscopy	ABSENT	-	Absent
RED BLOOD CELLS	Microscopy	2.5	cells/HPF	0-5
URINARY LEUCOCYTES (PUS CELLS)	Microscopy	2.5	cells/HPF	0-5
EPITHELIAL CELLS	Microscopy	2.5	cells/HPF	0-5
CASTS	Microscopy	ABSENT	-	Absent
CRYSTALS	Microscopy	ABSENT	-	Absent
BACTERIA	Microscopy	ABSENT	-	Absent
YEAST	Microscopy	ABSENT	-	Absent
PARASITE	Microscopy	ABSENT	-	Absent

(Reference : *PEI - Protein error of indicator, *GOD-POD - Glucose oxidase-peroxidase)

Sample Collected on (SCT) : Sample collection time
 Sample Received on (SRT) : Sample receiving time at Lab
 Report Released on (RRT) : Report release time
 Sample Type : URINE
 Labode :
 Barcode :

Doctor 1 Sign

Doctor 2 Sign

PROCESSED AT :

Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400 703 | 9870666333 | wellness@thyrocare.com

First National Diagnostic Chain to have 100% of its Labs with NABL Accreditation[#]

NAME : XXXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXXX
TEST ASKED : BLOOD SUGAR (F)

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
FASTING BLOOD SUGAR(GLUCOSE)	PHOTOMETRY	90.1	mg/dL

Bio. Ref. Interval. :-

As per ADA Guideline: Fasting Plasma Glucose (FPG)	
Normal	70 to 100 mg/dl
Prediabetes	100 mg/dl to 125 mg/dl
Diabetes	126 mg/dl or higher

Note :

The assay could be affected mildly and may result in anomalous values if serum samples have heterophilic antibodies, hemolyzed, icteric or lipemic. The concentration of Glucose in a given specimen may vary due to differences in assay methods, calibration and reagent specificity. For diagnostic purposes results should always be assessed in conjunction with patients medical history, clinical findings and other findings.

Please correlate with clinical conditions.

Method:- GOD-PAP METHOD

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : FLUORIDE PLASMA
Labcode :
Barcode :

Doctor 1 Sign

Doctor 2 Sign

First National Diagnostic Chain to have 100% of its Labs with NABL Accreditation[#]

NAME : XXXXXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXXXXX
TEST ASKED : ELEMENTS 22 (TOXIC AND NUTRIENTS), HBA PROFILE, HEMOGRAM
SAMPLE COLLECTED AT : XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS	Bio. Ref. Interval.
ARSENIC	ICP-MS	< 2.46	µg/L	< 5
CADMIUM	ICP-MS	< 0.75	µg/L	< 1.5
MERCURY	ICP-MS	< 2.46	µg/L	< 5
LEAD	ICP-MS	< 74.95	µg/L	< 150
CHROMIUM	ICP-MS	< 15	µg/L	< 30
BARIUM	ICP-MS	< 14.96	µg/L	< 30
COBALT	ICP-MS	< 0.8	µg/L	0.10 - 1.50
CAESIUM	ICP-MS	< 2.46	µg/L	< 5
THALLIUM	ICP-MS	< 0.5	µg/L	< 1
URANIUM	ICP-MS	< 0.5	µg/L	< 1
STRONTIUM	ICP-MS	< 23	µg/L	8 - 38
ANTIMONY	ICP-MS	< 9.05	µg/L	0.10 - 18
TIN	ICP-MS	< 1	µg/L	< 2
MOLYBDENUM	ICP-MS	< 2.35	µg/L	0.70 - 4.0
SILVER	ICP-MS	< 2	µg/L	< 4
VANADIUM	ICP-MS	< 0.4	µg/L	< 0.8
BERYLLIUM	ICP-MS	< 0.45	µg/L	0.10 - 0.80
BISMUTH	ICP-MS	< 0.45	µg/L	0.10 - 0.80
SELENIUM	ICP-MS	< 200	µg/L	60 - 340
ALUMINIUM	ICP-MS	< 15	µg/L	< 30
NICKEL	ICP-MS	< 7.5	µg/L	< 15
MANGANESE	ICP-MS	< 13.55	µg/L	7.10 - 20

Please correlate with clinical conditions.

Method :

ICP - MASS SPECTROMETRY

Note: Reference range has been obtained after considering 95% population as cutoff.

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : EDTA Whole Blood
Labcode :
Barcode :

Doctor 1 Sign

Doctor 2 Sign

PROCESSED AT :



Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400 703 9870666333 wellness@thyrocare.com

First National Diagnostic Chain to have 100% of its Labs with NABL Accreditation[#]

NAME : XXXXXXXXXXXXXXXXXX

SAMPLE COLLECTED AT :

REF. BY : XXXXXXXXXXXXXXXXXX

XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST ASKED : ELEMENTS 22 (TOXIC AND NUTRIENTS),HBA
PROFILE,HEMOGRAM

TEST NAME	TECHNOLOGY	VALUE	UNITS
HbA1c - (HPLC)	H.P.L.C	5.1	%

Bio. Ref. Interval. :

Bio. Ref. Interval.: As per ADA Guidelines

Below 5.7% : Normal
5.7% - 6.4% : Prediabetic
>=6.5% : Diabetic

Guidance For Known Diabetics

Below 6.5% : Good Control
6.5% - 7% : Fair Control
7.0% - 8% : Unsatisfactory Control
>8% : Poor Control

Method : Fully Automated H.P.L.C method

AVERAGE BLOOD GLUCOSE (ABG) CALCULATED 100 mg/dL

Bio. Ref. Interval. :

90 - 120 mg/dl : Good Control
121 - 150 mg/dl : Fair Control
151 - 180 mg/dl : Unsatisfactory Control
> 180 mg/dl : Poor Control

Method : Derived from HBA1c values

Please correlate with clinical conditions.

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : EDTA Whole Blood
Labcode :
Barcode :

Doctor 1 Sign

Doctor 2 Sign

NAME : XXXXXXXXXXXXXXXXX

SAMPLE COLLECTED AT :

REF. BY : XXXXXXXXXXXXXXXXX

XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST ASKED : ELEMENTS 22 (TOXIC AND NUTRIENTS),HBA
PROFILE,HEMOGRAM

TEST NAME	METHODOLOGY	VALUE	UNITS	Bio. Ref. Interval.
HEMOGLOBIN	SLS-Hemoglobin Method	10.4	g/dL	13.0-17.0
Hematocrit (PCV)	CPH Detection	33.6	%	40.0-50.0
Total RBC	HF & EI	3.88	X 10⁶/μL	4.5-5.5
Mean Corpuscular Volume (MCV)	Calculated	86.6	fL	83.0-101.0
Mean Corpuscular Hemoglobin (MCH)	Calculated	26.8	pq	27.0-32.0
Mean Corp.Hemo. Conc (MCHC)	Calculated	31	g/dL	31.5-34.5
Red Cell Distribution Width - SD (RDW-SD)	Calculated	48.9	fL	39-46
Red Cell Distribution Width (RDW - CV)	Calculated	15.4	%	11.6-14
RED CELL DISTRIBUTION WIDTH INDEX (RDWI)	Calculated	343.7	-	*Refer Note below
MENTZER INDEX	Calculated	22.3	-	*Refer Note below
TOTAL LEUCOCYTE COUNT (WBC)	HF & FC	7.77	X 10³ / μL	4.0 - 10.0
DIFFERENTIAL LEUCOCYTE COUNT				
Neutrophils Percentage	Flow Cytometry	75.2	%	40-80
Lymphocytes Percentage	Flow Cytometry	17.4	%	20-40
Monocytes Percentage	Flow Cytometry	2.7	%	2-10
Eosinophils Percentage	Flow Cytometry	3.9	%	1-6
Basophils Percentage	Flow Cytometry	0.5	%	0-2
Immature Granulocyte Percentage (IG%)	Flow Cytometry	0.3	%	0-0.5
Nucleated Red Blood Cells %	Flow Cytometry	0.1	%	0.0-5.0
ABSOLUTE LEUCOCYTE COUNT				
Neutrophils - Absolute Count	Calculated	5.84	X 10 ³ / μL	2.0-7.0
Lymphocytes - Absolute Count	Calculated	1.35	X 10 ³ / μL	1.0-3.0
Monocytes - Absolute Count	Calculated	0.21	X 10 ³ / μL	0.2 - 1.0
Basophils - Absolute Count	Calculated	0.04	X 10 ³ / μL	0.02 - 0.1
Eosinophils - Absolute Count	Calculated	0.3	X 10 ³ / μL	0.02 - 0.5
Immature Granulocytes (IG)	Calculated	0.02	X 10 ³ / μL	0-0.3
Nucleated Red Blood Cells	Calculated	0.01	X 10 ³ / μL	0.0-0.5
PLATELET COUNT	HF & EI	260	X 10³ / μL	150-410
Mean Platelet Volume (MPV)	Calculated	10.3	fL	6.5-12
Platelet Distribution Width (PDW)	Calculated	11.2	fL	9.6-15.2
Platelet to Large Cell Ratio (PLCR)	Calculated	26.6	%	19.7-42.4
Plateletcrit (PCT)	Calculated	0.27	%	0.19-0.39

*Note - Mentzer index (MI), RDW-CV and RDWI are hematological indices to differentiate between Iron Deficiency Anemia (IDA) and Beta Thalassemia Trait (BTT). MI >13, RDWI >220 and RDW-CV >14 more likely to be IDA. MI <13, RDWI <220, and RDW-CV <14 more likely to be BTT. Suggested Clinical correlation. BTT to be confirmed with HB electrophoresis if clinically indicated.

Method : Fully automated bidirectional analyser (6 Part Differential SYSMEX XN-1000)

(Reference : *FC- flowcytometry, *HF- hydrodynamic focussing, *EI- Electric Impedence, *Hb- hemoglobin, *CPH- Cumulative pulse height)

-- End of report --

Sample Collected on (SCT) : Sample collection time
 Sample Received on (SRT) : Sample receiving time at Lab
 Report Released on (RRT) : Report release time
 Sample Type : EDTA Whole Blood
 Labcode :
 Barcode :

Doctor 1 Sign

Doctor 2 Sign

CONDITIONS OF REPORTING

- v The reported results are for information and interpretation of the referring doctor only.
- v It is presumed that the tests performed on the specimen belong to the patient; named or identified.
- v Results of tests may vary from laboratory to laboratory and also in some parameters from time to time for the same patient.
- v Should the results indicate an unexpected abnormality, the same should be reconfirmed.
- v Only such medical professionals who understand reporting units, reference ranges and limitations of technologies should interpret results.
- v This report is not valid for medico-legal purpose.
- v Neither Thyrocare, nor its employees/representatives assume: (a) any liability, responsibility for any loss or damage that may be incurred by any person as a result of presuming the meaning or contents of the report, (b) any claims of any nature whatsoever arising from or relating to the performance of the requested tests as well as any claim for indirect, incidental or consequential damages. The total liability, in any case, of Thyrocare shall not exceed the total amount of invoice for the services provided and paid for.
- v Thyrocare Discovery video link :- <https://youtu.be/nbdYeRgYyQc>

EXPLANATIONS

- v Majority of the specimen processed in the laboratory are collected by Pathologists and Hospitals we call them as "Clients".
- v **Name** - The name is as declared by the client and recored by the personnel who collected the specimen.
- v **Ref.Dr** - The name of the doctor who has recommended testing as declared by the client.
- v **Labcode** - This is the accession number in our laboratory and it helps us in archiving and retrieving the data.
- v **Barcode** - This is the specimen identity number and it states that the results are for the specimen bearing the barcode (irrespective of the name).
- v **SCP** - Specimen Collection Point - This is the location where the blood or specimen was collected as declared by the client.
- v **SCT** - Specimen Collection Time - The time when specimen was collected as declared by the client.
- v **SRT** - Specimen Receiving Time - This time when the specimen reached our laboratory.
- v **RRT** - Report Releasing Time - The time when our pathologist has released the values for Reporting.
- v **Reference Range** - Means the range of values in which 95% of the normal population would fall.

SUGGESTIONS

- v Values out of reference range requires reconfirmation before starting any medical treatment.
- v Retesting is needed if you suspect any quality shortcomings.
- v Testing or retesting should be done in accredited laboratories.
- v For suggestions, complaints, clinical support or feedback, write to us at customersupport@thyrocare.com or call us on **022-309 0000**

Preventive Healthcare is now at your fingertips!



Explore & Select Test / Profile



Book Through App



Booking Confirmation



Track your Technician



Blood Collection



Sample Testing



Download Report & Reorder

Download Thyroapp Now




— Launching —

Jaanh

For a closer look at your health with

Doctor-Curated Specialised Packages All Under One Roof

 Heart	 Fever	 Cancer	 Women's - Reproductive Health
 Thyroid	 Diabetes	 STDs	 Skin Care
		 Hair Fall	



Thyrocare
Tests you can trust

+T&C Apply, #As on 5th December 2024, *As per a survey on doctors' perception of laboratory diagnostics (IJARIIT,2023)